



LYME DISEASE IN PENNSYLVANIA

A Supplement to the Report Issued by the

Task Force on Lyme Disease and Related Tick-Borne Diseases

Pursuant to Act 83 of 2014

September 2015

LYME DISEASE REFERENCES ONLY

Lyme disease

Adrion et al. Health Care Costs, Utilization and Patterns of Care following Lyme Disease. PLOSOne. 2015.
<http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0116767>

Aguero-Rosenfeld, M. E. (2008). Lyme disease: laboratory issues. *Infect. Dis. Clin. North Am.* 22, 301–313. doi: 10.1016/j.idc.2007.12.005 [PubMed Abstract](#) | [PubMed Full Text](#) | [CrossRef Full Text](#)

Allen, Herbert B.*, Hannaway, Maria; Joshi, Suresh; "Tertiary Treponematosis" Department of Dermatology, Drexel University College of Medicine, Philadelphia, Pennsylvania *Corresponding author: Herbert B. Allen, Department of Dermatology, Drexel University College of Medicine, Philadelphia, Pennsylvania, Jun 30, 2015, *J Clin Exp Dermatol Res* 2015, 6:4, <http://dx.doi.org/10.4172/2155-9554.10000288>.

Ang, C. W., Notermans, D. W., Hommes, M., Simoons-Smit, A. M., and Herremans T. (2011). Large differences between test strategies for the detection of anti-Borrelia antibodies are revealed by comparing eight ELISAs and five immunoblots. *Eur. J. Clin. Microbiol. Infect. Dis.* 30, 1027–1032. doi: 10.1007/s10096-011-1157-6. [PubMed Abstract](#) | [PubMed Full Text](#) | [CrossRef Full Text](#)

Arroja et al. Lyme carditis. *Rev Med Suisse.* 2015 Mar 4;11(464):563-6, 568.
<http://www.ncbi.nlm.nih.gov/pubmed/25924252>

Assous, M. V. (2007). Laboratory methods for the diagnosis of clinical forms of Lyme borreliosis (in French). *Med. Mal. Infect.* 37, 487–495. doi: 10.1016/j.medmal.2006.01.019 [PubMed Abstract](#) | [PubMed Full Text](#) | [CrossRef Full Text](#)

Aucott, J. N., A. W. Rebman, et al. (2013). "Post-treatment Lyme disease syndrome symptomatology and the impact on life functioning: is there something here?" *Qual Life Res* 22(1): 75-84.

Auwaerter, P. G., Bakken, J. S., Dattwyler, R. J., Dumler, J. S., Halperin, J. J., McSweeney, E., et al. (2011).

Antiscience and ethical concerns associated with advocacy of Lyme disease. *Lancet Infect. Dis.* 11, 713–719. doi: 10.1016/S1473-3099(11)70034-2. [PubMed Abstract](#) | [PubMed Full Text](#) | [CrossRef Full Text](#)

Bacon et al. Serodiagnosis of Lyme disease by kinetic enzyme-linked immunosorbent assay using recombinant VlsE1 or peptide antigens of Borrelia burgdorferi compared with 2-tiered testing using whole-cell lysates. *JID* 2003, 187: 1187 <http://jid.oxfordjournals.org/content/187/8/1187.long>

Baker, PJ. Pathogenesis of Lyme Neuroborreliosis in an Animal Model of Infection. *American Journal of Pathology* Vol 185 No 8, August 2015. [http://ajp.amjpathol.org/article/S0002-9440\(15\)00278-3/abstract](http://ajp.amjpathol.org/article/S0002-9440(15)00278-3/abstract)

Bakken LL, Callister SM, Wand PJ, Schell RF. Interlaboratory comparison of test results for detection of Lyme disease by 516 participants in the Wisconsin State Laboratory of Hygiene/College of American Pathologists Proficiency Testing Program. *J. Clin. Microbiol.* 35(3), 537–543 (1997).

Barthold SW, Hodzic E, Imai DM, Feng S, Yang X, Luft BJ. Ineffectiveness of tigecycline against persistent Borrelia burgdorferi. *Antimicrobial agents and chemotherapy.* Feb;54(2):643-51.

Belum et al. The Jarische-Herxheimer reaction: Revisited. *Travel Med Infect Dis.* 2013 Jul-Aug; 11(4):231

Bennet, R., Lindgren, V., and Zwegyberg-Wirgart, B. (2008). Borrelia antibodies in children evaluated for Lyme neuroborreliosis. *Infection* 36, 463–466. doi: 10.1007/s15010-008-6259-4 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Betheja et al. *J Neuropsychiatry Clin Neurosci.* 2013 Feb 27. Post-treatment Lyme Syndrome and central sensitization:<http://neuro.psychiatryonline.org/doi/full/10.1176/appi.neuropsych.12090223>

Blanc, F., Jaulhac, B., Fleury, M., de Sèze, J., de Martino, S. J., Rémy, V., et al. (2007). Relevance of the antibody index to diagnose Lyme neuroborreliosis among seropositive patients. *Neurology* 69, 953–958. doi: 10.1212/01.wnl.0000269672.17807.e0. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Branda et al. 2-tiered antibody testing for early and late Lyme disease using only an immunoglobulin G blot with the addition of a VlsE band as the second-tier test. *CID* 2010, 50:2

Branda, J. A., and Rosenberg, E. (2013). S. *Borrelia miyamotoi*: a lesson in disease discovery. *Ann. Intern. Med.* 159, 61–62. doi: 10.7326/0003-4819-159-1-201307020-00009 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Bransfield RC. Preventable cases of autism: relationship between chronic infectious diseases and neurological outcome. *Pediatric Health.* (2009) April 3(2).

Brorson, O., and Brorson, S. H. (1998). In vitro conversion of *Borrelia burgdorferi* to cystic forms in spinal fluid, and transformation to mobile spirochetes by incubation in BSK-H medium. *Infection* 26, 144–150. doi: 10.1007/BF02771839. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Brunner, M. (2006). Report refuting value of immune complexes to diagnose Lyme disease is invalid. *Clin. Vaccine Immunol.* 13, 304–306. doi: 10.1128/0142-1122.13.2.304-306.2006 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Burgess EC. *Borrelia burgdorferi* infection in Wisconsin horses and cows. *Ann N Y Acad Sci.* 1988;539:235-43. <http://www.ncbi.nlm.nih.gov/pubmed/3190095>

Cameron et al. Evidence assessments and guideline recommendations in Lyme disease: the clinical management of known tick bites, erythema migrans rashes and persistent disease. *Expert Rev. Anti Infect. Ther.* 1-33 (2014). <http://www.ncbi.nlm.nih.gov/pubmed/25077519>

Cameron et al. Proof that Chronic Lyme Disease exists. *Interdisciplinary Perspectives on Infectious Disease* (2010). <http://www.hindawi.com/journals/ipid/2010/876450/>

Carr, R. Pontiac Tribune. Chronic Lyme Disease: A New Era of Ignorance. March 14, 2015. <http://pontiactribune.com/chronic-lyme-disease-new-era-ignorance/>

Caskey et al. Persistence development by *B. burgdorferi* Populations in Vitro. *Antimicrob. Agents Chemother.* 27 July 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26248368>

CDC. (2011). Centers for Disease Control and Prevention. National notifiable Diseases Surveillance System. Lyme disease - *Borrelia burgdorferi* - 2011 Case Definition. Available online at: www.cdc.gov/NNDSS/script/casedef.aspx (Accessed March 24, 2014).

CDC. (2013). Centers for Disease Control and Prevention. Media Relations. CDC Provides Estimate of Americans Diagnosed with Lyme Disease Each Year. Available online at: www.cdc.gov/media/releases/2013/p0819-lyme-disease.html (Accessed March 24, 2014).

Ceftriaxone-associated biliary complications of treatment of suspected disseminated Lyme disease--New Jersey, 1990-1992. *MMWR Morb Mortal Wkly Rep.* 1993 Jan 22;42(2):39-42.
<http://www.cdc.gov/mmwr/preview/mmwrhtml/00018860.htm>

Centers for Disease Control and Prevention, Lyme disease: Two-step Laboratory Testing Process
<http://www.cdc.gov/lyme/diagnosis/testing/LabTest/TwoStep/index.html>

Centers for Disease Control and Prevention. Notice to Readers: Caution Regarding Testing for Lyme Disease. *Mmwr.* February 11, 2005 February 11, 2005;54((05)):125.

Chancellor et al. Urinary dysfunction in Lyme disease. *J Urol.* 1993 Jan;149(1):26-30.
<http://www.ncbi.nlm.nih.gov/pubmed/8417211>

Chiao et al. Antigens of Lyme disease of spirochaete *Borrelia burgdorferi* inhibits antigen or mitogen-induced lymphocyte proliferation. *FEMS Immunol Med Microbiol.* 1994 Feb;8(2):151-5.
<http://www.ncbi.nlm.nih.gov/pubmed/8173554>

Chmielewska-Badora, J., Cisak, E., Wojcik-Fatla, A., Zwolinski, J., Buczek, A., and Dutkiewicz, J. (2006). Correlation of tests for detection of *Borrelia burgdorferi sensu lato* infection in patients with diagnosed borreliosis. *Ann. Agric. Environ. Med.* 13, 307–311. [Pubmed Abstract](#) | [Pubmed Full Text](#)

Chmielewski et al. Improvement in the laboratory recognition of lyme borreliosis with the combination of culture and PCR methods. *Mol Diagn.* 2003;7(3-4):155-62. <http://www.ncbi.nlm.nih.gov/pubmed/15068385>

Clark, K. L., Leydet, B., and Hartman, S. (2013). Lyme borreliosis in human patients in Florida and Georgia, USA. *Int. J. Med. Sci.* 10, 915–931. doi: 10.7150/ijms.6273. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Connecticut Attorney General's Office. Attorney General's Investigation Reveals Flawed Lyme Disease Guideline Process, IDSA Agrees To Reassess Guidelines, Install Independent Arbiter, May 1, 2008.
<http://www.ct.gov/ag/cwp/view.asp?A=2795&Q=414284>

Cook et al. Lyme borreliosis: a review of data on transmission time after tick attachment. *Int J Gen Med.* 2014 Dec 19;8:1-8.
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4278789/>

Corona A. 1. *Borrelia burgdorferi*: Carbon Metabolism and the Tick-Mammal Enzootic Cycle. *Microbiol Spectr.* 2015 Jun; 3(3). <http://www.ncbi.nlm.nih.gov/pubmed/26185064>

Cottle et al. Lyme disease in a British referral clinic. *QJM*. 2012 Jun;105(6):537.
<http://qjmed.oxfordjournals.org/content/105/6/537.long>

Coulter, P, Lema, C et al; Two-Year Evaluation of *Borrelia burgdorferi* Culture and Supplemental Tests for Definitive Diagnosis of Lyme Disease, *Journal of Clinical Microbiology*, October 2005, p. 5080-5084, Vol. 43, No. 10.

Delong, A. K., B. Blossom, et al. (2012). "Antibiotic retreatment of Lyme disease in patients with persistent symptoms: A biostatistical review of randomized, placebo-controlled, clinical trials." *Contemp Clin Trials*.

Dietrich, T., Geissdörfer, W., Schlötzer-Schrehardt, U., Holbach, L., Schoerner, C., and Seitz, B. (2008). *Borrelia*-associated crystalline keratopathy with intracorneal detection of *Borrelia garinii* by electron microscopy and polymerase chain reaction. *Cornea* 27, 498–500. doi: 10.1097/ICO.0b013e318162a8f5. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Djukic M, Schmidt-Samoa C, Nau R, von Steinbüchel N, Eiffert H, Schmidt H., The diagnostic spectrum in patients with suspected chronic Lyme neuroborreliosis--the experience from one year of a university hospital's Lyme neuroborreliosis outpatients clinic. *Eur J Neurol*. 2011 Apr;18(4):547-55. doi: 10.1111/j.1468-1331.2010.03229.x. Epub 2010 Oct 27.

Donta ST. Late and chronic Lyme disease. *Med. Clin. North Am.* 86(2), 341–349 (2002).

Donta, S. T., Noto, R. B., and Vento, J. A. (2012). SPECT brain imaging in chronic Lyme disease. *Clin. Nucl Med* 37, 219–222. doi: 10.1097/RLU.0b013e318262ad9b. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Dorward et al. Immune capture and detection of *Borrelia burgdorferi* antigens in urine, blood, or tissues from infected ticks, mice, dogs, and humans. *J Clin Microbiol*. 1991 Jun;29(6):1162-70.
<http://www.ncbi.nlm.nih.gov/pubmed/1864935>

Eisendle et al. The expanding spectrum of cutaneous borreliosis. *G Ital Dermatol Venereol*. 2009 Apr;144(2):157-71.
<http://www.ncbi.nlm.nih.gov/pubmed/19357623>

Elsner et al. Suppression of Long-Lived Humoral Immunity Following *Borrelia burgdorferi* Infection. *PLoSOne*. 2015.
<http://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1004976>

Embers, M. E., Barthold, S. W., Borda, J. T., Bowers, L., Doyle, L., Hodzic, E., et al. (2012). Persistence of *Borrelia burgdorferi* in Rhesus macaques following antibiotic treatment of disseminated infection. *PLoS ONE* 7:e29914. doi: 10.1371/journal.pone.0029914. Erratum *PLoS ONE* 7. doi: 10.1371. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Engstrom SM, Shoop E, Johnson RC. Immunoblot interpretation criteria for serodiagnosis of early Lyme disease. *J. Clin. Microbiol.* 33(2), 419–427 (1995). Fallon et al. A Comparison of Lyme Disease Serologic Test Results From 4 Laboratories in Patients With Persistent Symptoms After Antibiotic Treatment. *Clin Infect Dis*. (2014) 59 (12): 1705-1710. <http://cid.oxfordjournals.org/content/59/12/1705.abstract>

EUCALB. (1997). European Concerted Action on Lyme Borreliosis, Diagnosis: Serology: Minimum Standards. Available online at: www.eucalb.com (Accessed March 24, 2014).

Fallon BA, Lipkin RB, Corbera KM et al: "Regional Cerebral Blood Flow and Metabolic Rate in Persistent Lyme Encephalopathy. Arch Gen Psychiatry. 2009;66(5):554-563.

Fallon et al. A Comparison of Lyme Disease Serologic Test Results From 4 Laboratories in Patients With Persistent Symptoms After Antibiotic Treatment. Clin Infect Dis. (2014) 59 (12): 1705-1710.
<http://cid.oxfordjournals.org/content/59/12/1705.abstract>

Fallon et al. A randomized, placebo-controlled trial of repeated IV antibiotic therapy for Lyme encephalopathy. Neurology 2008. 70(13):992-1003. <http://www.neurology.org/content/70/13/992.long>

Feder et al. A Critical Appraisal of "Chronic Lyme Disease". NEMJ 2007;357:1422-30.
<http://www.nejm.org/doi/pdf/10.1056/NEJMra072023>

Fehr, J. S., Bloemberg, G. V., Ritter, C., Hombach, M., Lüscher, T. F., Weber, R., et al. (2010). Septicemia caused by tick-borne bacterial pathogen *Candidatus Neohrlichia mikurensis*. Emerging Infect. Dis. 16, 1127–1129. doi: 10.3201/eid1607.091907. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Feng, Jie, and Megan Weitner, Wanliang Shi, Shuo Zhang, David Sullivan and Ying Zhang, Identification of Additional Anti-Persister Activity against *Borrelia burgdorferi* from an FDA Drug Library, Antibiotics 2015, 4(3), 397-410. <http://dx.doi.org/10.3390/antibiotics4030397>

Gallo et al. Amyloid-DNA Composites of Bacterial Biofilms Stimulate Autoimmunity. Immunity 42, 1171–1184, June 16, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26084027>

Gerber et al. Lyme disease in children in southeastern Connecticut. Pediatric Lyme Disease Study Group. N Engl J Med. 1996 Oct 24;335(17):1270-4. <http://www.nejm.org/doi/full/10.1056/NEJM199610243351703>

Halperin et al. Common Misconceptions About Lyme Disease. Amer J of Med, Vol 126, No 3, March 2013):
<http://www.amjmed.com/article/S0002-9343%2812%2900875-3/pdf>

Han GS et al. Exposure to *Borrelia burgdorferi* and other tick-borne pathogens in Gettysburg National Military Park, South-Central Pennsylvania, 2009. Vector Borne Zoonotic Dis. 2014 Apr;14(4):227-33.

Hao, Q., Hou, X., Geng, Z., and Wan, K. (2011). Distribution of *Borrelia burgdorferi sensu lato* in China. J. Clin. Microbiol. 49, 647–650. doi: 10.1128/JCM.00725-10. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Hodzic, E., Feng, S., Holden, K., Freet, K. J., and Barthold, S. W. (2008). Persistence of *Borrelia burgdorferi* following antibiotic treatment in mice. Antimicrob. Agents Chemother. 52, 1728–1736. doi: 10.1128/AAC.01050-07. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Holl-Wieden, A., Suerbaum, S., and Girschick, H. J. (2007). Seronegative Lyme arthritis. Rheumatol. Int. 27, 1091–1093. doi: 10.1007/s00296-007-0333-6. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Hoogman et al. Cognitive outcomes in adults following bacterial meningitis. J Neurol Neurosurg Psychiatry. 2007;78:1092-1096 <http://jnnp.bmj.com/content/78/10/1092.long>

Horowitz, MD, Richard, *Why Can't I Get Better? Solving the Mystery of Lyme and Chronic Disease* (New York: St. Martin's Press, 2013)

Hunfeld, K. P., Ruzic-Sabljić, E., Norris, D. E., Kraiczky, P., and Strl, F. (2005). In vitro susceptibility testing of *Borrelia burgdorferi* sensu lato isolates cultured from patients with erythema migrans before and after antimicrobial chemotherapy. *Antimicrob. Agents Chemother.* 49, 1294–1301. doi: 10.1128/AAC.49.4.1294-1301.2005. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Hutchinson et al. Prevalence Rates of *Borrelia burgdorferi* (Spirochaetales: Spirochaetaceae), *Anaplasma phagocytophilum* (Rickettsiales: Anaplasmataceae), and *Babesia microti* (Piroplasmida: Babesiidae) in Host-Seeking *Ixodes scapularis* (Acari: Ixodidae) From Pennsylvania. *J Med Ent.* E-published ahead of print. <http://jme.oxfordjournals.org/content/jmedent/early/2015/04/14/jme.tjv037.full.pdf?ijkey=uHxEa0KCPB8ki3l&keytype=ref>

Johnson et al. Assessment of new culture method for detection of *Borrelia* species from serum of Lyme disease patients. *J Clin Microbiol.* 2014 Mar;52(3):721-4. <http://jcm.asm.org/content/52/3/721.long>.
Comment: <http://jcm.asm.org/content/52/5/1803.long> ; reply: <http://jcm.asm.org/content/52/5/1804.long>

Johnson et al. Healthcare access and burden of care for patients with Lyme disease: A large United States survey. September 2011. [http://www.healthpolicyjrnl.com/article/S0168-8510\(11\)00101-1/abstract?cc=y](http://www.healthpolicyjrnl.com/article/S0168-8510(11)00101-1/abstract?cc=y)

Johnson, L., Wilcox, S., Mankoff, J. and Stricker, RB (2014) Severity of Chronic Lyme Disease Compared to Other Chronic Conditions: A Quality of Life Survey. *PeerJ*, DOI 10.7717/peerj.322. (Open access.)

Kalish et al. Persistence of Immunoglobulin M or Immunoglobulin G Antibody Responses to *Borrelia burgdorferi*. 10–20 Years after Active Lyme Disease. *Clin Infect Dis.* (2001) 33 (6): 780-785.

Keilp JG, Corbera KM, Slavov I et al: "WAIS-III and WMS-III performance in chronic Lyme disease *J International Neuropsychological Society.* 2006;12(1):119-129.

Keller, A., Graefen, A., Ball, M., Matzas, M., Boisguerin, V., Maixner, F., et al. (2012). New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. *Nat. Commun.* 3:698. doi: 10.1038/ncomms1701. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Kelly, J. Lyme Culture Test Causes Uproar. *Medscape* (Jan 30, 2013).

Kennedy, A. G. (2013). Differential diagnosis and the suspension of judgment. *J. Med. Philos.* 38, 487–500. doi: 10.1093/jmp/jht043. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Kern et al. Heterogeneity of *Borrelia burgdorferi* Sensu Stricto Population and Its Involvement in *Borrelia* Pathogenicity: Study on Murine Model with Specific Emphasis on the Skin Interface. <http://www.ncbi.nlm.nih.gov/pubmed/26197047>

Klempner et al. Two controlled trials of antibiotic treatment in patients with persistent symptoms and a history of Lyme disease. 2001 *NEJM*, 345. <http://www.nejm.org/doi/full/10.1056/NEJM200107123450202>

Klempner et al. Intralaboratory reliability of serologic and urine testing for Lyme disease *Am J of Med*, Volume 110, Issue 3, 15 February 2001, Pages 217-219, Comment and Reply: Volume 111, Issue 6, 15 October 2001, Pages 502-503.

Klempner et al. Treatment Trials for Post-Lyme Disease Symptoms Revisited. *Am J Med*. 2013 Aug; 126(8): 665–669. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4075423/?report=reader>. Comments and replies: *Am J Med*. 2014 Feb; 127(2):e7-e12.

Krupp et al. Study and treatment of post Lyme disease (STOP-LD): a randomized double masked clinical trial. 2003, *Neurology*, 60. <http://www.neurology.org/content/60/12/1923.long>

Kubánek M et al. Detection of *Borrelia burgdorferi* sensu lato in endomyocardial biopsy specimens in individuals with recent-onset dilated cardiomyopathy. 1. *Eur J Heart Fail*. 2012 Jun;14(6):588-96. <http://www.ncbi.nlm.nih.gov/pubmed/22379178>

Kuchynka P et al. A high frequency of viral agents yet absence of *Borrelia burgdorferi* is seen within the myocardium of subjects with normal left ventricular systolic function: an electron microscopy study. *Folia Microbiol (Praha)*. 2015 Jul 24. <http://www.ncbi.nlm.nih.gov/pubmed/26205424>

Kuchynka P et al. Recent-onset dilated cardiomyopathy associated with *Borrelia burgdorferi* infection. *Herz*. 2015 May 5. <http://www.ncbi.nlm.nih.gov/pubmed/25939436>

Lee, D. J., and Vielmeyer, O. (2011). Analysis of overall level of evidence behind infectious diseases society of america practice guidelines. *Arch. Intern. Med*. 171, 18–22. doi: 10.1001/archinternmed.2010.482. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Lee, S. H., Vigliotti, J. S., Vigliotti, V. S., Jones, W., and Shearer, D. M. (2014). Detection of *Borreliae* in archived sera from patients with clinically suspect Lyme disease. *Int. J. Mol. Sci*. 15, 4284–4298. doi: 10.3390/ijms15034284. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Lewis, K. (2007). Persister cells, dormancy and infectious disease. *Nature* 5, 48–56. doi: 10.1038/nrmicro1557 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Liveris et al. Comparison of five diagnostic modalities for direct detection of *Borrelia burgdorferi* in patients with early Lyme disease. *Diagn Microbiol Infect Dis*. 2012 July ; 73(3): 243–245. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3377843/pdf/nihms368616.pdf>

Logigian, E. L., R. F. Kaplan, et al. (1990) “Chronic neurologic manifestations of Lyme disease.” *N Engl J Med* 323, 1438-1444.

Lopes de Carvalho, I., Fonseca, J. E., Marques, J. G., Ullmann, A., Hojgaard, A., Zeidner, N., et al. (2009). Vasculitis-like syndrome associated with *Borrelia lusitaniae* infection. *Clin. Rheumatol*. 27, 1587–1591. doi: 10.1007/s10067-008-1012-z. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Ma B, Christen B, Leung D, Vigo-Pelfrey C. Serodiagnosis of Lyme borreliosis by western immunoblot: reactivity of various significant antibodies against *Borrelia burgdorferi*. *J. Clin. Microbiol*.30(2), 370–376 (1992).

MacDonald A. Gestational Lyme borreliosis. Implications for the fetus. *Rheum Dis Clin North Am*. 1989 Nov;15(4):657-77.

Magri JM1, Johnson MT, Herring TA, Greenblatt JF. Lyme disease knowledge, beliefs, and practices of New Hampshire primary care physicians. *J Am Board Fam Pract*. 2002 Jul-Aug;15(4):277-84.

Makhani, N., Morris, S. K., Page, A. V., Brophy, J., Lindsay, L. R., Banwell, B. L., et al. (2011). A twist on Lyme: the challenge of diagnosing European Lyme neuroborreliosis. *J. Clin. Microbiol.* 49, 455–457. doi: 10.1128/JCM.01584-10. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Malloy et al. *Clin Infect Dis*. False-Positive Results of PCR Testing for Lyme Disease. (2001) 33 (3): 412-413. <http://cid.oxfordjournals.org/content/33/3/412.long>

Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases, 8th Edition. 2015.

Mann, L. Chicago Tribune. Lyme disease still stuck in controversy, but there is hope. August 4, 2015. <http://www.chicagotribune.com/lifestyles/health/sc-hlth-0805-lyme-disease-hope-20150730-story.html#page=1>

Mantovani, E., Costa, I. P., Gauditano, G., Bonoldi, V. L. N., Higuchi, M. L., and Yoshinari, N. H. (2007). Description of Lyme disease-like syndrome in Brazil. Is it a new tick borne disease or Lyme disease variation? *Braz. J. Med. Biol. Res.* 40, 443–456. doi: 10.1590/S0100-879X2006005000082. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Marangoni, A., Sparacino, M., Cavrini, F., Storni, E., Mondardini, V., Sambri, V., et al. (2005). Comparative evaluation of three different ELISA methods for the diagnosis of early culture-confirmed Lyme disease in Italy. *J. Med. Microbiol.* 54, 361–367. doi: 10.1099/jmm.0.45853-0. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Margos, G., Piesman, J., Lane, R. S., Ogden, N. H., Sing, A., Straubinger, R. K., et al. (2014). *Borrelia kurtenbachii* sp. nov.: a widely distributed member of the *Borrelia burgdorferi* sensu lato species complex in North America. *Int. J. Syst. Evol. Microbiol.* 64, 128–130. doi: 10.1099/ijms.0.054593-0. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Marques et al. Natural killer cell counts are not different between patients with post-Lyme disease syndrome and controls. *Clin Vaccine Immunol*. 2009 Aug;16(8):1249-50. <http://cvi.asm.org/content/16/8/1249.full.pdf+html>

Marques et al. Xenodiagnosis to detect *Borrelia burgdorferi* infection: a first-in-human study. *Clin Infect Dis*. 2014 Apr;58(7):937-45. <http://cid.oxfordjournals.org/content/58/7/937.long>

Mavin, S., Evans, R., Milner, R. M., Chatterton, J. M. W., and Ho-Yen, D. O. (2009). Local *Borrelia burgdorferi* sensu stricto and *Borrelia afzelii* strains in a single mixed antigen improves Western blot sensitivity. *J. Clin. Pathol.* 62, 552–554. doi: 10.1136/jcp.2008.063461. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Mavin, S., Milner, R. M., Evans, R., Chatterton, J. M. W., Joss, A. W. L., and Ho-Yen, D. O. (2007). The use of local isolates in Western blots improves serological diagnosis of Lyme disease in Scotland. *J. Med. Microbiol.* 56, 47–51. doi: 10.1099/jmm.0.46793-0. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Middelveen et al. Exploring the association between Morgellons disease and Lyme disease: identification of *Borrelia burgdorferi* in Morgellons disease patients. *BMC Dermatol.* 2015 Feb 12;15:1.
<http://www.ncbi.nlm.nih.gov/pubmed/25879673>

Miklossy J (2012) Chronic or late Lyme neuroborreliosis: analysis of evidence compared to chronic or late neurosyphilis. *Open Neurol J* 6: 146-157. Nelson et al. Neoplasms Misdiagnosed as “Chronic Lyme Disease”. *JAMA Intern Med.* 2015;175(1):132-133. <http://archinte.jamanetwork.com/article.aspx?articleid=1921752>

Miklossy J, Gern L, Darekar P, Janzer RC, Loos H. Senile plaques, neurofibrillary tangles and neuropil threads contain DNA? *J Spirochetal and Tick-borne Dis* 1995; 2: 1-5.

Miklossy J, Kasas S, Zurn AD, McCall S, Yu S, and McGeer PL. Persisting atypical and cystic forms of *Borrelia burgdorferi* and local inflammation in Lyme neuroborreliosis. *J Neuroinflammation* 2008; 5: 40-57.

Miklossy J, Kris A, Radenovic A, Miller L, Forro L, Martins R, Reiss K, Darbinian N, Darekara P, Mihaly L, and Khalili K. Beta amyloid deposition and Alzheimer’s type changes induced by *Borrelia spirochetes*. *Neurobiol Aging* 2006; 27: 228-236.

Miklossy J, Rosemberg S, and McGeer PL; Beta amyloid deposition in the atrophic form of general paresis. In *Alzheimer’s Disease: New advances.* Medimond. Proceedings of the 10th International Congress on Alzheimer’s Disease. Edited by: Iqbal K, Winblad B, and Avila J; 2006: 429-433.

Miklossy J. Chronic inflammation and amyloidogenesis in Alzheimer’s disease — role of spirochetes. *J Alzheimers Dis* 2008; 13: 381-391.

Miklossy JM, Khalili K, Gern L, Ericson RL, Darekar P, Bolle L, Hurlimann J, and Paster BJ. *Borrelia burgdorferi* persists in the brain in chronic Lyme neuroborreliosis and may be associated with Alzheimer’s disease. *J Alzheimers Dis* 2004; 6: 639-649.

Miklossy, J. 2008. Biology and neuropathology of dementia in syphilis and Lyme disease. In *Handbook of Clinical Neurology*, Vol. 89. C. Duyckaerts, I. Litvan (eds.). Elsevier, Amsterdam, Netherlands. p. 825-844. [Persistence of *B. burgdorferi* is evident in dementia patients.]

Miklossy, J. 2011. Alzheimer’s disease – a neurospirochetosis. Analysis of the evidence following Koch’s and Hill’s criteria. 2011; 8: 90 (<http://www.jneuroinflammation.com/content/8/1/90>) [91% of the brains of Alzheimer’s patients sampled were positive for spirochetes; 25% of Alzheimer’s patients analyzed had *B. burgdorferi* spirochetes in their brains. Persistence occurs when spirochetes change physical characteristics by converting to dormant cysts.]

Moyer M. The growing global battle against blood-sucking ticks. *Nature News* Vol 524, 27 August 2015.
<http://www.nature.com/news/the-growing-global-battle-against-blood-sucking-ticks-1.18227>

Murgia, R., and Cinco, M. (2004). Induction of cystic forms by different stress conditions in *Borrelia burgdorferi*. *APMIS* 112, 57–62. doi: 10.1111/j.1600-0463.2004.apm1120110.x. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Nelson et al. Neoplasms Misdiagnosed as “Chronic Lyme Disease”. JAMA Intern Med. 2015;175(1):132-133. <http://archinte.jamanetwork.com/article.aspx?articleid=1921752>

Nicolle, C. (1993). “Destin des maladies infectieuses (in French),” in Reedited by the Association des ed France Lafayette (Paris: Anciens Elèves de l'Institut Pasteur), 216.

Nields, J. A., and Kveton, J. F. (1991). Tullio phenomenon and seronegative Lyme borreliosis. Lancet 338, 128–129. doi: 10.1016/0140-6736(91)90130-H. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Nowakowski et al. Laboratory diagnostic techniques for patients with early Lyme disease associated with erythema migrans: a comparison of different techniques. Clin Infect Dis. (2001) 33 (12): 2023-2027. <http://cid.oxfordjournals.org/content/33/12/2023.long>

O’Day and Catalano. A Lack of Correlation between the Incidence of Lyme Disease and Deaths due to Alzheimer’s Disease. Journal of Alzheimer’s Disease 42 (2014) 115–118. <http://iospress.metapress.com/content/a96r5pw3714p617m/fulltext.pdf>

Ogden, N. H., Margos, G., Aanensen, D. M., Drebot, M. A., Feil, E. J., Hanincova, K., et al. (2011). Investigation of genotypes of *Borrelia burgdorferi* in *Ixodes scapularis* ticks collected during surveillance in Canada. Appl. Environ. Microbiol. 77, 3244–3254. doi: 10.1128/AEM.02636-10 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Pachner et al. Inoculation of nonhuman primates with the N40 strain of *Borrelia burgdorferi* leads to a model of Lyme neuroborreliosis faithful to the human disease. Neurology. 1995 Jan;45(1):165-72.

Palecek et al. Presence of *Borrelia burgdorferi* in endomyocardial biopsies in patients with new-onset unexplained dilated cardiomyopathy. Med Microbiol Immunol. 2010 May;199(2):139-43. <http://www.ncbi.nlm.nih.gov/pubmed/20052487>

Patel et al. Death from Inappropriate Therapy for Lyme Disease Clin Infect Dis. (2000) 31 (4): 1107-1109. <http://cid.oxfordjournals.org/content/31/4/1107.full>

Peer Reviewed Evidence of Persistence of Lyme Disease Spirochete *Borrelia burgdorferi* and Tick-Borne Diseases. http://www.ilads.org/ilads_news/2015/list-of-700-articles-citing-chronic-infection-associated-with-tick-borne-disease-compiled-by-dr-robert-bransfield/

Perronne C (2014) Lyme and associated tick-borne diseases: global challenges in the context of a public health threat. Front. Cell. Infect. Microbiol. 4:74. doi: 10.3389/fcimb.2014.00074.

Perronne, C. (2012). Lyme disease antiscience. Lancet Infect. Dis. 12, 361–362. doi: 10.1016/S1473-3099(12)70053-1. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Pfeiffer, Mary Beth, Exclusive: Dicey Path to Lyme Disease Diagnosis, The Poughkeepsie Journal, October 2014, <http://www.poughkeepsiejournal.com/story/news/health/lyme-disease/2014/03/21/lyme-diagnosis-false-tests/6692353/>

Pfister, H. W., V. Preac-Mursic, et al. (1991). "Randomized comparison of ceftriaxone and cefotaxime in Lyme neuroborreliosis." *J Infect Dis* 163(2): 311-318.

Phillips, S. E., Mattman, L. H., Hulinska, D., and Moayad, H. (1998). A proposal for the reliable culture of *Borrelia burgdorferi* from patients with chronic Lyme disease, even from those previously aggressively treated. *Infection* 26, 364–367. doi: 10.1007/BF02770837. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Qiu W-G, Bruno JF, McCaig WD, Xu Y, Livey I, Schriefer ME, et al. Wide distribution of a high-virulence *Borrelia burgdorferi* clone in Europe and North America. *Emerg Infect Dis* [serial on the Internet]. 2008 Jul [date cited]. Available from <http://wwwnc.cdc.gov/eid/article/14/7/07-0880>

Qureshi, M, Bedlack, RS, and Cudkowicz, ME. Lyme disease serology in amyotrophic lateral sclerosis. *Muscle and Nerve* 40: 626-628, 2009.

Ramesh et al. Inflammation in the pathogenesis of lyme neuroborreliosis, *AM J Pathol* 2015 May;185(5):1344-60. <http://www.ncbi.nlm.nih.gov/pubmed/25892509> Earlier work: *J Neuroinflammation*. 2013 Jul 18; <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3721987/>

Rizzoli, A., Hauffe, H. C., Carpi, G., Vourc'h, G. I., Neteler, M., and Rosà R. (2011). Lyme borreliosis in Europe. *Eurosurveillance* 16, 1–8. [Pubmed Abstract](#) | [Pubmed Full Text](#)

Roche-Lanquetot, M. O., Ader, F., Durand, M. C., Carlier, R., Defferière, H., Dinh, A., et al. (2008). Results of a prospective standardized study of 30 patients with chronic neurological and cognitive disorders after tick bites (in French). *Med. Mal. Infect.* 38, 543–548. doi: 10.1016/j.medmal.2008.06.007. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Rudenko, N., Golovchenko, M., and Grubhoffer, L., Oliver, J. H. Jr. (2011). Updates on *Borrelia burgdorferi* sensu lato complex with respect to public health. *Ticks Tick Borne Dis.* 2, 123–128. doi: 10.1016/j.ttbdis.2011.04.002. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Rudenko, N., Golovchenko, M., Ruzek, D., Piskunova, N., Mallatova, N., and Grubhoffer, L. (2009). Molecular detection of *Borrelia bissettii* DNA in serum samples from patients in the Czech Republic with suspected borreliosis. *FEMS Microbiol. Let.* 292, 274–281. doi: 10.1111/j.1574-6968.2009.01498.x. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Sapi E, Pabbati N, Datar A, Davies EM, Rattelle A, Kuo BA. Improved culture conditions for the growth and detection of borrelia from human serum. *Int J Med Sci.* 2013;10(4):362-76.

Schutzer SE1, Fraser-Liggett CM, Casjens SR, Qiu WG, Dunn JJ, Mongodin EF, Luft BJ. Whole-genome sequences of thirteen isolates of *Borrelia burgdorferi*. *J Bacteriol.* 2011 Feb;193(4):1018-20. doi: 10.1128/JB.01158-10. *Epub* 2010 Oct 8.

Schutzer, S. E., Coyle, P. K., Belman, A. L., Golightly, M. G., and Drulle, J. (1990). Sequestration of antibody to *Borrelia burgdorferi* in immune complexes in seronegative Lyme disease. *Lancet* 335, 312–315. doi: 10.1016/0140-6736(90)90606-6 [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Schutzer, Steven E., Angel, Thomas E.; Liu, Tao; Schepmoes, et al. "Distinct Cerebrospinal Fluid Proteomes Differentiate Post-Treatment Lyme Disease from Chronic Fatigue Syndrome", PLoS ONE, February 23, 2011. •DOI: 10.1371/journal.pone.0017287

Schwan et al. Interstitial cystitis and *Borrelia burgdorferi*. *Ann Intern Med*. 1989 Sep 15;111(6):537.
<http://annals.org/article.aspx?articleid=703382>

Schwan, T. G., Anderson, J. M., Lopez, J. E., Fischer, R. J., Raffel, S. J., McCoy, B. N., et al. (2012). Endemic foci of the tick-borne relapsing fever spirochete *Borrelia crocidurae* in Mali, West Africa, and the potential for human infection. *PLoS Negl. Trop. Dis.* 6:e1924. doi: 10.1371/journal.pntd.0001924. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Seinost G1, Golde WT, Berger BW, Dunn JJ, Qiu D, Dunkin DS, Dykhuizen DE, Luft BJ, Dattwyler RJ. Infection with multiple strains of *Borrelia burgdorferi sensu stricto* in patients with Lyme disease. *Arch Dermatol*. 1999 Nov;135(11):1329-33.

Seltzer et al. Long-term Outcomes of Persons With Lyme Disease. *JAMA*. 2000;283(5):609-616.
<http://jama.jamanetwork.com/article.aspx?articleid=192358>

Seriburi et al. High frequency of false positive IgM immunoblots for *Borrelia burgdorferi* in clinical practice. *Clin Microbiol Infect* 2012; 18: pp. 1236-1240

Shadick, N. A., C. B. Phillips, et al. (1999). "Musculoskeletal and neurologic outcomes in patients with previously treated Lyme disease." *Ann Intern Med* 131(12): 919-926.

Shah et al. Improved Sensitivity of Lyme disease Western Blots Prepared with a Mixture of *Borrelia Burgdorferi* Strains 297 and B31. *Chronic Diseases - International*. October 10 2014. <http://austinpublishinggroup.com/chronic-diseases/download.php?file=fulltext/chronicdiseases-v1-id1009.pdf>

Sharma et al. *Borrelia burgdorferi*, the causative agent of Lyme disease, forms 2 drug-tolerant persister cells. *Antimicrob. Agents Chemother.* (2015)

Sigal LH. Summary of the first 100 patients seen at a Lyme disease referral center. *Am J Med* 1990;88:577-81.
<http://www.ncbi.nlm.nih.gov/pubmed/2346158>

Skogman et al. Long-term Clinical Outcome After Lyme Neuroborreliosis in Childhood. *Pediatrics* Vol. 130 No. 2 August 1, 2012. pp. 262 -269.

Stanek, G., and Reiter, M. (2011). The expanding Lyme *Borrelia* complex—clinical significance of genomic species? *Clin. Microbiol. Infect.* 17, 487–493. doi: 10.1111/j.1469-0691.2011.03492.x. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Steere et al. The overdiagnosis of Lyme disease. *JAMA* 1993;269:1812-26.
<http://www.ncbi.nlm.nih.gov/pubmed/8459513>

Steere et al. Treatment of the early manifestations of Lyme. *Annals of Internal Medicine*. 1983. 99:22.

Steere et al. Prospective Study of Serologic Tests for Lyme Disease. CID 2008;47 (15 July).
<http://cid.oxfordjournals.org/content/47/2/188.full.pdf+html>

Stephens et al. Intralaboratory Reliability of Serologic and Urine Testing for Lyme Disease. American Journal of Medicine Vol 111, 502-503. <http://www.ncbi.nlm.nih.gov/pubmed/11690582>

Straubinger, R. K. (2000). PCR-based quantification of *Borrelia burgdorferi* organisms in canine tissues over a 500-day postinfection period. J. Clin. Microbiol. 38, 2191–2199. Pubmed Abstract | Pubmed Full Text

Straubinger, R. K., Summers, B. A., Chang, Y. F., and Appel, M. J. (1997). Persistence of *Borrelia burgdorferi* in experimentally infected dogs after antibiotic treatment. J. Clin. Microbiol. 35, 111–116. Pubmed Abstract | Pubmed Full Text

Stricker et al. Sexual transmission of Lyme disease: challenging the tickborne disease paradigm. Expert Rev. Anti Infect. Ther. Early online, 1–4 (2015).
<http://www.tandfonline.com/doi/abs/10.1586/14787210.2015.1081056?journalCode=ierz20#.VfsNAhFViko>

Stricker RB, Johnson L. Lyme wars: let's tackle the testing. BMJ. 2007 Nov 17;335(7628):1008.

Stricker, R. B., and Johnson, L. (2014). Lyme disease: call for a “Manhattan Project” to combat the epidemic. PLoS Pathog. 10:e1003796. doi: 10.1371/journal.ppat.1003796. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Stricker, Raphael B. and Lorraine Johnson, Lyme Disease: the next decade, Infect Drug Resist. 2011; 4: 1–9. Published online 2011 Jan 7. doi: 10.2147/IDR.S15653

Strle K, Jones KL, Drouin EE, Li X, Steere AC. *Borrelia burgdorferi* RST1 (OspC Type A) Genotype Is Associated with Greater Inflammation and More Severe Lyme Disease. Am J Pathol. 2011 Jun;178(6):2726-39.

Strle, F., and Stanek, G. (2009). Clinical manifestations of Lyme borreliosis. Curr. Probl. Dermatol. 37, 51–110. doi: 10.1159/000213070. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

Tager, F. A., Fallon, B. A., Keilp, J., Rissenberg, M., and Ray Jones, C., Liebowitz, M. R. (2001). A controlled study of cognitive deficits in children with chronic Lyme disease. J. Neuropsychiatry Clin. Neurosci. 13, 500–507. doi: 10.1176/appi.neuropsych.13.4.500. Pubmed Abstract | Pubmed Full Text | CrossRef Full Text

The Clinical Assessment, Treatment, and Prevention of Lyme Disease, Human Granulocytic Anaplasmosis, and Babesiosis: Clinical Practice Guidelines by the Infectious Diseases Society of America:
<http://cid.oxfordjournals.org/content/43/9/1089.full>

Tijssse-Klasen, Ellen, and Marion P. G. Koopmans, and Hein Sprong, Tick-borne pathogen - reversed and conventional discovery of disease, Front. Public Health, 07 July 2014 <http://dx.doi.org/10.3389/fpubh.2014.00073>,
1Centre for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven,
2Erasmus Medical Center, Rotterdam, Netherlands.

Timmaraju et al. Biofilm formation by 1 *Borrelia sensu lato*. FEMS Microbiology Letters, July 24, 2015. <http://femsle.oxfordjournals.org/content/362/15/fnv120.full>

Treib, J., A. Fernandez, et al. (1998). "Clinical and serologic follow-up in patients with neuroborreliosis." *Neurology* 51(5): 1489-1491.

Tveitnes, D., Oymar, K., and Natas, O. (2009). Laboratory data in children with Lyme neuroborreliosis, relation to clinical presentation and duration of symptoms. *Scand. J. Infect. Dis.* 41, 355–362. doi: 10.1080/00365540902787666. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Valesova, H., J. Mailer, et al. (1996). "Long-term results in patients with Lyme arthritis following treatment with ceftriaxone." *Infection* 24(1): 98-102.

Varela, A. S., Luttrell, M. P., Howerth, E. W., Moore, V. A., Davidson, W. R., Stallknecht, D. E., et al. (2004). First culture isolation of *Borrelia lonestari*, putative agent of Southern tick-associated rash illness. *J. Clin. Microbiol.* 42:1163–1169. doi: 10.1128/JCM.42.3.1163-1169.2004. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Vayssier-Taussat, M., Moutailler, S., Michelet, L., Devillers, E., Bonnet, S., Cheval, J., et al. (2013). Next generation sequencing uncovers unexpected bacterial pathogens in ticks in Western Europe. *PLoS ONE* 8:e81439. doi: 10.1371/journal.pone.0081439. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Wallet, F., Labalette, P., Herwegh, S., Loiez, C., Margaron, F., and Courcol, R. J. (2008). Molecular diagnosis of a bilateral panuveitis due to *Borrelia burgdorferi sensu lato* by cerebral spinal fluid analysis. *J. Infect. Dis.* 61, 214–215. [Pubmed Abstract](#) | [Pubmed Full Text](#)

Winjgaard et al. The burden of Lyme borreliosis expressed in disability-adjusted life years. *European Journal of Public Health* 1-8, June 16, 2015. <http://www.ncbi.nlm.nih.gov/pubmed/26082446>

Wojciechowska-Koszko, I., Maczynska, I., Szych, Z., and Giedrys-Kalemba, S. (2011). Serodiagnosis of borreliosis: indirect immunofluorescence assay, enzyme-linked immunosorbent assay and immunoblotting. *Arch Immunol. Ther. Exp.* 59, 69–77. doi: 10.1007/s00005-010-0111-0. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Wormser, G. P. (2006). Early Lyme disease. *N. Engl. J. Med.* 354, 2794–2801. doi: 10.1056/NEJMcp061181. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Wormser, G. P., Dattwyler, R. J., Shapiro, E. D., Halperin, J. J., Steere, A. C., Klempner, M. S., et al. (2006). The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin. Infect. Dis.* 43, 1089–1134. doi: 10.1086/508667. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Yrjänäinen, H., Hytönen, J., Hartiala, P., Oksi, J., and Viljanen, M. K. (2010). Persistence of borrelial DNA in the joints of *Borrelia burgdorferi*-infected mice after ceftriaxone treatment. *APMIS* 118, 665–673. doi: 10.1111/j.1600-0463.2010.02615.x. [Pubmed Abstract](#) | [Pubmed Full Text](#) | [CrossRef Full Text](#)

Zhang et al. Economic impact of Lyme Disease. *Emerging Infectious Diseases* Vol. 12 No 4 April 2006. pp. 653-660. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3294685/>

Ziska MH1, Donta ST, Demarest FC. Physician preferences in the diagnosis and treatment of Lyme disease in the United States. *Infection*. 1996 Mar-Apr;24(2):182-6.